

NEW EXPRESSIONS FOR LAGUERRE AND HERMITE POLYNOMIALS

H. MOYA-CESSA

ABSTRACT. New expressions for Laguerre and Hermite polynomials are shown. They are based on operator algebras commonly used in quantum mechanics.

1. HERMITE POLYNOMIALS

Hermite functions are given by the expression [1]

$$(1.1) \quad H_n(x) = (-1)^n e^{x^2} \frac{d^n}{dx^n} e^{-x^2}$$

by doing $p = -id/dx$, we may rewrite them as

$$(1.2) \quad H_n(x) = (-i)^n e^{x^2} p^n e^{-x^2}$$

given that $[x, p]f(x) = if(x)$ and considering the relation $e^{\mu A} B e^{-\mu A} = B + \mu[A, B] + \mu^2/2![A, [A, B]] + \dots$ we can cast (1.1) as

$$(1.3) \quad H_n(x) = (-i)^n (p + 2ix)^n.$$

We can add the Hermite polynomials given in (1.3) to obtain

$$(1.4) \quad \sum_{n=0}^{\infty} H_n(x) \frac{\alpha^n}{n!} = e^{-i\alpha(p+ix)}$$

by applying the Baker-Hausdorff formula [2], $e^{A+B} = e^{-[A,B]/2} e^A e^B$, valid when the commutators obey $[A, [A, B]] = [B, [A, B]] = 0$, we obtain

$$(1.5) \quad \sum_{n=0}^{\infty} H_n(x) \frac{\alpha^n}{n!} = e^{-\alpha^2 + 2\alpha x} e^{-i\alpha p} = e^{-\alpha^2 + 2\alpha x}$$

1.1. Addition formula. We want to apply the form obtained in (1.3) to evaluate the quantity

$$H_n(x+y),$$

we write it as

$$(1.6) \quad H_n(x+y) = (-i)^n \left[-i \frac{d}{d(x+y)} + 2i(x+y) \right]^n$$

by using the chain rule we have

$$(1.7) \quad \frac{d}{d(x+y)} = \frac{1}{2} \left(\frac{\partial}{\partial x} + \frac{\partial}{\partial y} \right)$$

Date: September 6th, 2008.

2000 Mathematics Subject Classification. Primary .

Key words and phrases. Polynomials, Operator algebras.

so that we may re-express (1.6) in the form

$$(1.8) \quad H_n(x+y) = \left(\frac{-i}{\sqrt{2}}\right)^n \left(-i\frac{\partial}{\partial\sqrt{2}x} + 2i\sqrt{2}x - i\frac{\partial}{\partial\sqrt{2}y} + 2i\sqrt{2}y\right)^n$$

by defining

$$(1.9) \quad p_X = -i\frac{\partial}{\partial X}, \quad p_Y = i\frac{\partial}{\partial Y}$$

with $X = \sqrt{2}x$ and $Y = \sqrt{2}y$ we obtain

$$(1.10) \quad H_n(x+y) = \frac{1}{2^{n/2}} \sum_{k=0}^n \binom{n}{k} (-i)^k (p_X + 2iX)^k \times (-i)^{n-k} (p_Y + 2iY)^{n-k}$$

that by using (1.3) adds to

$$(1.11) \quad H_n(x+y) = \frac{1}{2^{n/2}} \sum_{k=0}^n \binom{n}{k} H_k(\sqrt{2}x) H_{n-k}(\sqrt{2}y)$$

2. LAGUERRE POLYNOMIALS

Laguerre polynomials are given by

$$(2.1) \quad L_n^\alpha(x) = \frac{1}{n!} x^{-\alpha} e^x \frac{d^n}{dx^n} (e^{-x} x^{n+\alpha})$$

that may be rewritten as

$$(2.2) \quad L_n^\alpha(x) = \frac{1}{n!} x^{-\alpha} e^x (ip)^n e^{-x} x^{n+\alpha}$$

by using that $e^x p^n e^{-x} = (p+i)^n$ we can cast them into

$$(2.3) \quad L_n^\alpha(x) = \frac{1}{n!} x^{-\alpha} \left(\frac{d}{dx} - 1\right)^n x^{n+\alpha}$$

or

$$(2.4) \quad L_n^\alpha(x) = \frac{1}{n!} x^{-\alpha} \sum_{m=0}^n \binom{n}{m} (-1)^{n-m} \frac{d^m}{dx^m} x^{n+\alpha}$$

because

$$(2.5) \quad \frac{d^m}{dx^m} x^{n+\alpha} = \frac{(n+\alpha)!}{(n+\alpha-m)!} x^{n+\alpha-m}$$

so we obtain the usual form for Laguerre polynomials:

$$(2.6) \quad L_n^\alpha(x) = \sum_{k=0}^n \binom{n+\alpha}{n-k} (-1)^k \frac{x^k}{k!}.$$

3. CONCLUSIONS

We have given new forms for Hermite polynomials (1.3) and Laguerre polynomials (2.3) that may be used to add these polynomials in an easier form.

REFERENCES

- [1] I.S. Gradshteyn, I.M. Ryzhik, Table of Integrals, Series and Products, Academic Press, Inc., San Diego, CA, 1980.
- [2] W.H. Louisell, Quantum Statistical Properties of Radiation, Wiley, New York, 1973.

INAOE, APDO. POSTAL 51 Y 216, 72000, PUEBLA, PUE., MEXICO

E-mail address: `hmmc@inaoe.mx`